

The role of Helmet use on Severity and Pattern of Maxillofacial Injuries among Motorcycle Crash Victims attending Mulago Hospital, Uganda

Joses O. Komakech¹, Isaac Okullo¹, Francis Lakor², Charles Mugisha Rwenyonyi^{1*}

1.Department of Dentistry, College of Health Sciences, Makerere University, Kampala

2.Oral and Maxillofacial Surgery Unit, Mulago Hospital, Kampala

Abstract

The objective of the study was to establish the role of helmet use on severity and pattern of maxillofacial injuries among motorcycle crash victims attending Mulago Hospital. This was descriptive cross sectional case series study using a questionnaire in form of an oral interview, clinical examination and review of medical records of the patients. The pattern of injuries was assessed based on the demarcated regions of the face. The severities of the injuries were scored using the Facial Injury Severity Scale. The data were analysed using SPSS version 17.0. There were 105 participants (male/female: 97/8) aged 15 to 63 years. Most participants (n= 74, 70.5%) were riders and the rest were passengers. There were no female riders. Most participants were from Kampala and surrounding districts. Overall, 35.2% of the participants had helmets: 50% of the riders and 0% of the passengers. About 73% of the participants used Three Quarter Shell helmet design. There was no significant difference in the pattern of injuries between the passengers and non helmeted riders ($p=0.076$). There was a higher frequency helmeted riders (n = 15, 40.5%) with pan-facial fractures compared to other participants. Non helmeted riders and passengers had significantly more injuries to the lower two thirds of the face compared to the helmeted riders. Passengers had less severe facial injuries compared to riders. We conclude that about a third of the participants who were riders used helmets, being particularly the Three Quarter Shell design. The helmeted riders had more severe maxillofacial injuries and of panfacial category as compared to other participants. The passengers had least severe injuries compared to riders. It is recommended to carry out further studies to elucidate on motorcycle related maxillofacial injuries especially in regard to the quality of helmets and their adequate use.

Corresponding author:-Charles Mugisha Rwenyonyi, Department of Dentistry College of Health Sciences, Makerere University, P.O.Box 7072, Kampala Uganda

e-mail: mrwenyonyi@chs.mak.ac.ug, Tel. +256772405349/+256751405349

Running title:- Role of crash helmet on severity and pattern of maxillofacial injuries

Keywords: FISS, helmet, maxillofacial injuries, motorcycle crash, severity

Received : 08 Jan 2017;

Accepted :25 Jun 2017;

Published :27 Jun 2017

Introduction

Motorcycles are a common means of transportation in low and middle income countries. Their use is associated with high prevalence of injuries; a major public health concern¹. Previous studies^{2,3} indicated a direct relationship between the frequency of motorcycle usage and injuries in different countries. Injuries due to motorcycle crashes are highest in South East Asian countries compared to other countries¹.

In Nigeria, maxillo-facial injuries due to motorcycle riding constituted 50.5% to 57% of the road traffic accidents^{4,5}. In East Africa, the trend of fatalities due to motorcycle crashes is much lower and varies with countries: 16%, Rwanda; 7%, Uganda and Tanzania, and 1%, Kenya⁶. Overall, injuries due to motorcycle crashes in Kampala metropolitan is estimated at about 25%⁷ and those involving the head region constitute about 35.5%⁸.

Kampala metropolitan is congested with public service vehicles; estimated at 15,000 minibuses and 30,000 motorcycles⁹. Currently, more motorcycles are being imported into Uganda^{6,10}, particularly for public transportation because of their ability to circumvent traffic congestion, versatility in reaching remote places inaccessible to public passenger vehicles and provision of employment to youths. Galukande et al.⁸ estimated that riding motorcycles for public transport provides employment to about 100,000 youths in Uganda. The role of helmets in protection against maxillofacial injuries is still conflicting. Some studies¹¹⁻¹³ have implicated helmets in exacerbating maxillofacial injuries (especially of the mandible), others^{14,15} reported that helmets protect the face against severe injuries, while others^{16,17} indicated that helmet use significantly reduced traumatic brain injuries. Although motorcycles provide quick public transport in Uganda, they are a risk to injury^{8,18}. A few studies^{8,18} on motorcycle crashes in Uganda reported general injuries of the body without specifying the maxillofacial injuries, while Kamulegeya et al.¹⁹ had a cursory information relating maxillofacial injuries and motorbikes/helmet use. The purpose of the present study was to investigate the role of helmet use in severity and pattern of maxillofacial injuries among motorcycle crash victims attending Mulago Hospital, Uganda

Material and Methods

Study design

This was case series descriptive cross sectional study.

Study setting

The study was carried out in Oral and Maxillofacial Surgery, Neurosurgery and Orthopaedics units of Mulago Hospital. Mulago Hospital is a national referral and teaching health facility located in Kampala, the capital city of Uganda. The hospital has a capacity of 1500 beds. The units are specialized in managing patients, particularly with trauma who are mainly referred from other health facilities in the country.

Selection of study participants

The study participants (n=105) were patients aged 15 – 63 years with maxillofacial injuries following motorcycle crashes either as riders or passengers and attending Oral and Maxillofacial Surgery (n=77), Neurosurgery (n=22) and Orthopaedic unit (n=6) of Mulago Hospital. They were requested for consent/assent before recruitment into the study. Participants who were too ill to answer the questionnaire (n=3) and too young to wear a helmet (n=4) were excluded from the study.

Data collection

A structured questionnaire was administered to the participants or their guardians in form of an oral interview by the Principal Investigator (JOK) to get background information. Information on the design of helmet worn during crash was obtained from the participants with the help of pictures of helmets (Fig. 1).

Clinical examination for maxillofacial injuries was carried out while the patient was lying supine in the dental chair/couch with the help of artificial light by JOK. The pattern of maxillofacial bone fracture was assessed clinically and confirmed with radiological imaging.. The severity of the injuries was scored on Facial Injury Severity Scale (FISS) and recorded in a data recording form.

Data management and analysis

Data on completed forms were double checked for errors and completeness while the participant was still present in the clinic or ward. The data were inputted



Fig. 1. The pictures of different helmet designs used by the riders in the present study

into a computer and again double checked for errors and completeness. The injury severity scores based on FISS were categorized: 1 = score 1-6 and 2 = score > 6. The data were analyzed using Statistical Package for Social Sciences Inc., (SPSS version 17.0 for windows, Chicago, Illinois, USA). Frequency distribution was used to summarize the data. Chi-square statistics was used to assess any significant differences in participants based on quantitative variables. Paired t test was used to assess the intra-examiner reliability in recording observations. The level of significance was set at p value <5%.

Reliability test

Blind duplicate examination of mandibular fractures was conducted on 20 patients 4 days after the main survey. The agreement was substantial (Cohen's $\kappa = 0.70$). Similarly, blind duplicate viewing of 50 radiographs, a week later gave an almost perfect agreement (Cohen's $\kappa = 0.84$). There was neither evidence of systematic error in clinical examination nor in viewing of radiographs (paired t test, $p > 0.05$).

Ethical Considerations

The study complied with a protocol approved by Makerere University School of Medicine Research and Ethics Committee (No. REC REF 2011-173). Permission to carry out the study was obtained from Mulago Hospital administration. Informed consent was obtained from the study participants /guardians of children below 18 years of age. In addition to the consent, the children were requested to assent to the study. The consent/assent form was translated from

English into the local language (Luganda) for participants who did not understand English. The nature and purpose of the study were explained to the participants in accordance with Helsinki Declaration²⁰. They were informed that they were at liberty to accept or refuse to participate in the study without being coerced and their refusal would not affect their relationship with investigators. No personal identifiers were used in recording the data and the final report on the findings is anonymous.

Results


The study sample constituted 105 participants (male/female: 97/8). Most participants ($n = 74$; 70.5%) were riders and none of them a female. They all rode the saddle motor cycles. Overall, the participants were aged 15 to 63 (mean, 29.01 ± 9.5) years with majority 62.9%, ($n = 66$) aged 21 to 30 years. The riders were aged 18 to 59 (mean, 28 ± 8.7) years while the passengers were 15 to 63 (mean, 29 ± 11.4) years. The helmeted riders were aged 18 to 59 years (mean, 30.4 ± 9.3) years and the non helmeted counterparts were 18 to 50 (mean, 27.1 ± 7.8) years. Based on the units of study, the helmeted riders were 22, 11 and 4 in Oral and maxillofacial, Neurosurgery and Orthopaedic Surgery unit, respectively.

Districts where maxillofacial injuries occurred

The subjects with maxillofacial injuries came from 21 districts of Uganda. Kampala district had the majority of cases (46.7%, $n = 49$) followed by Wakiso district with 21.0% ($n = 22$) and Mokono district had

((n=10, 9.5%; Fig. 2).

face compared to the helmeted riders ($p = 0.020$). Only



1 rider and with a helmet had injuries of the upper third of the face. The relationship between the pattern of

Helmet design and usage

Three different types of helmet design were used by the riders: the TQS ($n=27, 73.0\%$), FF ($n=8, 21.6\%$) and HS ($n=2, 5.4\%$). Overall, 35.2% of the participants had helmets: 50% of the riders and 0% of the passengers (Table 1-3). Older riders were more likely to wear helmets as compared with the younger counterpart (Table. 1).

There was no significant difference in the pattern of injuries between the passengers and non helmeted riders ($p=0.076$). There was a higher frequency of helmeted riders ($n = 15, 40.5\%$) with pan-facial fractures compared to 4 non helmeted riders and 1 passenger (Table 2). Non helmeted riders and passengers had significantly more injuries to the lower two thirds of the

maxillofacial injuries and helmet designs could not be assessed because of the skewed variability of helmet design.

Severity of maxillofacial injuries based on FISS

The influence of gender on FISS score could not be assessed because the data were skewed towards males. The overall mean FISS was 7.0 (SD 5.3). Helmeted riders had more severe maxillofacial injuries, range 1 to 19 (mean, 10.1 ± 5.4) FISS as compared to their non helmeted counterparts (mean FISS, 6.59 ± 4.9). Passen-

Table 1. The frequency distribution of participants according to helmet usage by age groups (n=105)

Age Groups		Helmet usage, n (%)		
		Helmeted	Non-helmeted	Total
< 21 years	Passenger	0 (0.0)	4 (100)	4 (100)
	Rider	2 (16.7)	10 (83.3)	12 (100)
≥21 years	Passenger	0 (0.0)	27 (100)	27 (100)
	Rider	35 (56.5)	27 (43.5)	62 (100)

Use of helmet significantly increased with age of the motorcycle riders (p=0.004). Pattern of maxillofacial injuries

Table 2. The frequency distribution of participants according to pattern of maxillofacial injuries and helmet use (n=105)

Helmet use		Injured part of the maxillofacial region, n (%)						
		UT	MT	LT	UTT	ULT	LTT	PF
Non helmeted	Passenger (n=31)	0 (0.0)	3 (9.7)	14 (45.2)	3 (9.7)	0 (0.0)	10 (32.3)	1 (3.2)
	Rider (n=37)	0 (0.0)	2 (5.4)	17 (45.9)	3 (8.1)	3 (8.1)	8 (21.6)	4 (10.4)
Helmeted	Rider (n=37)	1 (2.7)	4 (10.8)	3 (8.1)	4 (10.8)	1 (2.7)	9 (24.3)	15 (40.5)

UT = Upper Third; MT = Middle Third; LT = Lower Third; UTT = Upper Two Thirds; ULT = Upper and Lower Thirds; LTT = Lower Two Thirds; PF = Pan Facial

Table 3. The frequency distribution of participants according to severity of maxillofacial injury based on Facial Injury Severity Scale and helmet use (n=105)

Helmet use		FISS score, n (%)	
		01-Jun	>6
Non helmeted	Passenger (n=31)	30 (96.8%)	1 (3.2%)
	Rider (n=37)	24 (64.9%)	13 (35.1%)
Helmeted	Rider (n=37)	12 (32.4%)	25 (67.6%)

gers had the least severity of maxillofacial injuries (mean FISS, 4.0 ± 1.6 ; Table 3).

Helmeted riders had significantly more severe

from distant districts may have been managed in regional referral health facilities³³.

In the present study, the motorcycle riders were



Fig. 3. The picture of a passenger who sustained severe injuries involving loss of a left eye

maxillofacial injuries as compared to the non helmeted counterparts ($p=0.004$) and overall, the riders sustained significantly more severe maxillofacial injuries as compared to the passengers ($p=0.001$). Three of the participants lost an eye each: 2 helmeted riders (5.4%) and a female passenger (3.2%, Fig. 3), particularly who were ejected some distance from the motorcycle to tarmac.

Discussion

There are several scoring systems, but with challenges related to their use as previously discussed²¹⁻³² though without reaching a consensus. In the present study, FISS was used to record severity of maxillofacial injury because of its reliability²¹ and easy to compute²². We observed most of the participants were from Kampala and the neighboring districts probably because the study site (Mulago Hospital) is within Kampala and in proximity of the surrounding districts. Furthermore, Kampala metropolitan has more motorcycles compared to other regions⁹, hence the high frequency of the injured. Additionally, although Mulago Hospital is a national referral health facility, some of the trauma patients

aged 18 to 59 (mean, 28 ± 8.7) years, which is consistent with other studies elsewhere^{4,5,8,16,34}, but higher than reported in Indianapolis, United States of America³⁵. The lowest age permissible for acquiring a motorcycle riding license varies in different countries because of varying socio-economic, political and cultural backgrounds. In Uganda, motorcycles are mainly used for public transport, a business mainly involving youths⁸, while in other countries they are used so much in sports, leisure and children (as young as 14 years) commuting to school^{36,37}.

We observed all the riders were males, who coincidentally, are the ones mostly involved in riding motorcycles for public transport in Uganda. This finding is similar to previous studies^{5,8,38} in Africa. Elsewhere in the world, there are variable proportions of female riders: 2.2% in Mersey and North Wales, UK³⁴; 24.6% in Athens, Greece³⁹; 30% in Indianapolis, USA³⁵; 11.9% in Hyderabad, India⁴⁰ and 24% in Kerala, India¹⁵.

The absence of female motorcycle riders in the present study and elsewhere in Africa may probably be

due to the type of motorcycles on the market, the straddle type which do not favor mounting by female and child riders unlike the scooters, which are more appealing and commonly used in recreational and sporting activities⁴¹.

In the present study, 50% of the riders used helmet (Table 1 and 2) which was higher than the 17.7%⁸ and 29%³³ earlier reported among patients attending Mulago Hospital. The older participants were more likely to put on helmet compared to the younger age group (Table 1) and this was similar to observation in Vietnam⁴². However, this was self motivated usage of helmet because although there is a law⁴³ on mandatory use, it is not adequately enforced in Uganda similar to observation in Nigeria⁴, and contrary to the United Kingdom where the law is enforced and usage rate is 93.1%³⁴.

Other reasons for not willing to wear helmet include: discomfort, obscuring of vision, disturbance of hearing, ruffling up hair style, cost or thefts of helmets while others said helmets are responsible for crashes³⁹. However, in the present study, we were not able to interview the participants on the opinion about helmet use.

In the present study, about three quarters (27/37) of the riders wore helmets of Three Quarter Shell design and because of the limited variability, the relationship between helmet design and type of maxillofacial injuries could not be assessed. However, amongst the non helmeted participants, the injuries were predominantly mandibular fractures and the associated soft tissue injuries while the helmeted group had panfacial fractures in addition to mandibular fractures. Overall, most participants had injuries of the lower two thirds of the face irrespective of helmet use (Table 2). Furthermore, it may also be assumed that the Three Quarter Shell helmets predominantly used by the study participants lack the chin bar, which leaves the lower two third of the face exposed to injury. However, Yates and colleagues¹² reported higher incidence of mandibular fractures amongst the riders wearing Full Face type of helmet (with chin bar) as compared to non helmeted counterparts.

We found significantly more helmeted riders had severe maxillofacial injuries (with 2 of them losing an eye each) as compared to the non helmeted counter-

parts (Table 3), in support of Gopalakrishna et al.¹¹ Gopirikan et al.¹⁴ and Jayadevan et al.¹⁵, but contrary to findings from Yates et al.¹². Earlier on, Brian et al.¹³ revealed no significant difference in severe injuries between the helmeted and nonhelmeted groups. Our finding could be due to poor quality⁴⁴ or improper use of helmets like the size and fastening of the straps. However, we were neither able to assess the quality of the helmets nor the adequacy of their use.

Overall, we found the riders had sustained significantly more severe maxillofacial injuries as compared to the passengers contrary to previous workers⁴⁵ who found no significant difference between the riders and passengers. Additionally, the passengers had the least severe maxillofacial injuries compared to the riders (Table 3). It could be postulated that the bodies of the riders could have shielded the passengers who ordinarily sit behind the riders. However, one female passenger lost an eye (Fig. 3) and it was an isolated case where she was ejected from the motorcycle to some distance on the tarmac. This is consistent with Gopalakrishna et al.¹¹ report that participants with maxillofacial injuries were those who were ejected from the motorcycle to some distance during the crash.

One previous study⁴⁶ that used FISS as a scoring system recorded the most severely injured subject with FISS score of 11, which was much lower than 19 seen in the present study, The difference may probably be due to factors like differences in quality of the roads and motorcycle overloading. We could not assess the influence of gender on FISS score because of skewed gender distribution among the study participants.

Conclusion

About a third of the participants and only riders used helmets particularly of the Three Quarter Shell design at the time of injury. None of the passengers had a helmet. Generally, helmeted riders had severe maxillofacial injuries and of panfacial category as compared to other participants.

Recommendation

Although the present study generated baseline data, it was daunted with various limitations indicative of further studies to elucidate on factors such as the quality of helmets and their adequate use as well as mechanism of maxillofacial injuries during motorcycle crashes.

Acknowledgements

The authors are grateful to the participants for their cooperation during the study. The staff members in the study units of Mulago Hospital were instrumental in mobilizing the study participants.

Conflict of interest

The authors declare that there was no conflict of interest in the course of carrying out this study.

References

1. World Health Organisation. World report on road traffic prevention, 2004. Accessed on 20.6.2016 from {<http://apps.who.int/iris/bitstream/10665/42871/1/9241562609.pdf>}
2. Dubois L, Leijdesdorff HA, Goslings JC, Lam PH, Bui HL et al. Motorcycle related maxillofacial trauma in southern Vietnam: need for preventive measurements. *Intern J Oral Maxillofac Surg.* 2009;38 (5):579 -84.
3. Maitree S. Road accidents: A common persistent problem. Keynote address on scientific conference on Road Safety On Four Continents, Bangkok, Thailand, 2007.
4. Oginni FO, Ugboko VI, Ogundipe O, Adegbeingbe BO. Motorcycle-related maxillofacial injuries among Nigerian intracity road users. *J Oral Maxillofac Surg.* 2006;64:56-62.
5. Solagberu BA, Ofoegbu CKP, Nasir AA, Ogundipe OK, Adekanye AO et al. Motorcycle injuries in a developing country and the vulnerability of riders, passengers, and pedestrians. *Injury Prev.* 2006;12:266-8.
6. Odero W. Motorcycle Injuries In: East Africa: Risk factors and prevention. Paper presented at the scientific meeting of the RTIRN regional workshop, Accra, Ghana, 2009.
7. Naddumba EK. A cross-sectional retrospective study of boda boda injuries at Mulago Hospital in Kampala -Uganda. Makerere University, 2007.
8. Galukande M, Jombwe J, Fualal J, Gakwaya A. Boda-boda injuries a health problem and a burden of disease in Uganda: a tertiary hospital survey. *East Centr Afr J Surg.* 2009;14(2):33-7.
9. Wandera GO. Urban traffic congestion in African cities – An over view of Kampala. Kampala: Government, 2010.
10. Sserwaniko F. Indian bikes topple Japanese, 2009. Accessed on 15.6.2016 from {<http://www.newvision.co.ug/D/8220/672752>}
11. Gopalakrishna G, Peek-Asa C, Kraus JF. Epidemiologic features of facial injuries among motorcyclists. *Ann Emerg Med.* 1998;32:425-30.
12. Yates JM, Dickenson AJ. Helmet use and maxillofacial injuries sustained following low speed motorcycle accidents. *Injury.* 2002;33:479-83.
13. Brian MW, Bobek S, Dierks, EJ, Bell RB, Potter J et al. Comparison of helmeted and un-helmeted motorcyclists and evaluation of maxillofacial injuries. *J Oral Maxillofac Surg.* 2006;64(9 Suppl):33-4.
14. Thota G. Review of Motorcycle Related Facial Injuries Pre-and Post-2003 Pennsylvania Helmet Law Modification From 1/1998 to 8/2008 at Allegheny General Hospital, Pittsburgh, PA. *J Oral Maxillofac Surg.* 2009 Sep 30;67(9):62-7.
15. Jayadevan S, Jayakumary M, Binoo D, Jeesha CH. Determinants of safety helmet use among motorcyclists in Kerala, India. *J Injury Viol Res.* 2010; 2 (1): 49-4.
16. Ankarath S, Giannoudis PV, Barlow I, Bellamy MC, Matthews S J et al. Injury patterns associated with mortality following motorcycle crashes. *Injury.* 2002;33:473-7.
17. Servadei F, Begliomini C, Gardini E. Effect of Italy's motorcycle helmet law on traumatic brain injuries. *Injury Prev.* 2003;9:257-60.
18. Naddumba EK. A Cross sectional retrospective study of Boda Boda injuries at Mulago Hospital in Kampala, Uganda. *East Centr Afr J Surg.* 2004;19(1):44-7.
19. Kamulegeya A, Lakor F, Kabenge K. Oral maxillofacial fractures seen at a Ugandan tertiary hospital: a six-month prospective study. *Clinics.* 2009; 64 (9):843-8.
20. World Medical Association. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *J Am Med Assoc.* 2013 Nov 27;310(20):2191-4.
21. Chawda MN, Hildebrand F, Pape HC, Giannoudis PV.

- Predicting outcome after multiple trauma: which scoring system? *Injury*. 2004;35:347–58.
22. Bagheri SC, Dierks EJ, Kademani D, Holmgren E, Bell RB et al. Application of a Facial Injury Severity Scale in Craniomaxillofacial Trauma. *J Oral Maxillofac Surg*. 2006;64:408-14
 23. Kirkpatrick JR, Youmans RL. TRAUMA INDEX: an aide in the evaluation of injured victims. *J Trauma Acute Care Surg*. 1971;11(8):711-4.
 24. Baker SP, O'Neill B, Haddon Jr. W, Long WB. The injury severity score: a method for describing patients with multiple injuries and evaluating emergency care. *J Trauma Acute Care Surg*. 1974;14(3):187–96.
 25. Baker SP, O'Neill B. Injury severity score: an update. *J Trauma Acute Care Surg*. 1976 Nov 1;16(11):882-5
 26. Champion HR, Sacco WJ, Lepper RL. An anatomic index of injury severity. *J Trauma Acute Care Surg*. 1980 Mar 1;20(3):197-202.
 27. Knaus WA, Draper EA, Wagner DP, Zimmerman JE. APACHE II: a severity of disease classification system. *Critical Care Med*. 1985;13:818-29.
 28. Baxt WG, Moody P. The differential survival of trauma patients. *J Trauma Acute Care Surgery*. 1987 Jun 1;27(6):602-6
 29. Champion HR, Copes WS, Sacco WJ. A new characterization of injury severity. *J Trauma*. 1990;30(5):539–45.
 30. Bazzoli G J, Madura KJ, Cooper GF. Progress in the development of trauma systems. I. United States. *J Am Med Assoc*. 1995;273:395–401.
 31. Brenneman FD, Boulanger BR, McLellan BA, Redelmeier DA. Measuring injury severity: time for a change? *J Trauma Acute Care Surg*. 1998 Apr 1;44(4):580-2.. 1998;44(4):580–2.
 32. Al West T, Rivara FP, Cummings P, Jurkovich GJ, Maier RV. Harborview assessment for risk of mortality: an improved measure of injury severity on the basis of ICD-9-CM. *J Trauma Acute Care Surg*. 2000 Sep 1;49(3):530-41.
 33. Naddumba EK. Musculoskeletal Trauma Services in Uganda. *Clin Orthopaedic Rel Res*. 2008;466, 2317–22.
 34. Sinha AK, Boot DA, Gorman DF, Teanby DN. Severe motorcycle injury in Mersey region and North Wales. *Injury*. 1995 Oct 31;26(8):543-5.
 35. Mangus RS, Simons CJ, Jacobson LE. Current helmet and protective equipment usage among previously injured ATV and motorcycle riders. *Injury Prev*. 2004;1056-8.
 36. Maw-Chang L, Wen-Ta C, Li-Tung C, Liu SC, Lin SH. Craniofacial injuries in unhelmeted riders of motorbikes. *Injury*. 1995;26 (7):467-70.
 37. Segui-Gomez M, Lopez-Valdes FJ. Recognizing the importance of injury in other policy forums: the case of motorcycle licensing policy in Spain. *Injury Prev*. 2007;13:429-30.
 38. Andrew A. Motorcycle related trauma in South Sudan: a cross sectional observational study. *South Sud Med J* 2010;2(4):7-9.
 39. Skalkidou A, Petridou E, Papadopoulos FC. Factors affecting motorcycle helmet use in the population of Greater Athens, Greece. *Injury Prev*. 1999;5:264-7.
 40. Michael CH, Barry LE, Davi, WT, Stephen EB. Etiology and prevention of craniomaxillofacial trauma. In Peter Ward Booth, Barry L Eppley and Rainer Schmelzeisen (Eds.), *Maxillofacial trauma and esthetic facial reconstruction* (pp. 3-19). Philadelphia, PA: Elsevier Science, 2003.
 41. Zalavras C, Nikolopoulou G, Essin D, Manjra N, Lewis E et al. Pediatric Fractures During Skateboarding, Roller Skating, and Scooter Riding. *Am J Sports Med*. 2005;33(4):568-73
 42. Hung DV, Stevenson MR, Ivers RQ. Prevalence of helmet use among motorcycle riders in Vietnam. *Injury Prev*. 2006;12:409–13.
 43. Parliament of Uganda. Traffic and Road Safety Act 1998, Cap. 361, Part VIII.
 44. Mau-Roung L. Kraus JF. A review of risk factors and patterns of motorcycle injuries. *Accid Anal Prev* 2009;41:710–22.
 45. Fitzharris M, Dandona R, Kumar GA, Dandona L. Crash characteristics and patterns of injury among hospitalized motorized two-wheeled vehicle users in urban India. *BMC Public Health*. 2009 Jan 12;9(1):1.

46. Oginni FO, Ajike SO, Obuekwe ON, Fasola O. A prospective multicenter study of injury profile, severity and risk factors in 221 motorcycle-injured Nigerian maxillofacial patients. *Traffic Injury Prev.* 2009;10(1):70-5.